spet

ACETAMINOPHEN-INDUCED HEPATOTOXICITY

LAURA P. JAMES, PHILIP R. MAYEUX, AND JACK A. HINSON

Departments of Pediatrics (L.P.J.) and Pharmacology and Toxicology (L.P.J., P.R.M., J.A.H.), University of Arkansas for Medical Sciences, Little Rock, Arkansas

(Received March 20, 2003; accepted June 25, 2003)

This article is available online at http://dmd.aspetjournals.org

ABSTRACT:

The analgesic acetaminophen causes a potentially fatal, hepatic centrilobular necrosis when taken in overdose. The initial phases of toxicity were described in Dr. Gillette's laboratory in the 1970s. These findings indicated that acetaminophen was metabolically activated by cytochrome P450 enzymes to a reactive metabolite that depleted glutathione (GSH) and covalently bound to protein. It was shown that repletion of GSH prevented the toxicity. This finding led to the development of the currently used antidote N-acetylcysteine. The reactive metabolite was subsequently identified to be N-acetyl-p-benzoquinone imine (NAPQI). Although covalent binding has been shown to be an excellent correlate of toxicity, a number of other events have been shown to occur and are likely important in the initiation and repair of toxicity. Recent data have shown that nitrated tyrosine residues as well as acetaminophen adducts occur in the necrotic cells following toxic doses of acetaminophen. Nitrotyrosine was postulated to be mediated by peroxynitrite, a reactive nitrogen species formed by the very rapid reaction of superoxide and nitric oxide (NO). Peroxynitrite is normally detoxified by GSH, which is depleted in acetaminophen toxicity. NO synthesis (serum nitrate plus nitrite) was dramatically increased following acetaminophen. In inducible nitric oxide synthase (iNOS) knockout mice, acetaminophen did not increase NO synthesis or tyrosine nitration; however, histological evidence indicated no difference in toxicity. Acetaminophen did not cause hepatic lipid peroxidation in wild-type mice but did cause lipid peroxidation in iNOS knockout mice. These data suggest that NO may play a role in controlling lipid peroxidation and that reactive nitrogen/oxygen species may be important in toxicity. The source of the superoxide has not been identified, but our recent finding that NADPH oxidase knockout mice were equally sensitive to acetaminophen and had equal nitration of tyrosine suggests that the superoxide is not from the activation of Kupffer cells. It was postulated that NAPQI-mediated mitochondrial injury may be the source of the superoxide. In addition, the significance of cytokines and chemokines in the development of toxicity and repair processes has been demonstrated by several recent studies. IL-1 β is increased early in acetaminophen toxicity and may be important in iNOS induction. Other cytokines, such as IL-10, macrophage inhibitory protein-2 (MIP-2), and monocyte chemoattractant protein-1 (MCP-1), appear to be involved in hepatocyte repair and the regulation of proinflammatory cytokines.

Downloaded from dmd.aspetjournals.org by guest on February 16, 2012

Overdoses of the analgesic and antipyretic acetaminophen represent one of the most common pharmaceutical product poisonings in the United States today (Litovitz et al., 2002). Although considered safe at therapeutic doses, in overdose, acetaminophen produces a centrilobular hepatic necrosis that can be fatal (Prescott, 1980). Whereas the initial biochemical and metabolic events that occur in the early stages of toxicity have been well described, the precise mechanisms of hepatocyte death are poorly understood. Necrosis is recognized as the mode of cell death and apoptosis has been ruled out (Lawson et al., 1999; Gujral et al., 2002).

This work was supported by National Institute of Diabetes and Digestive and Kidney Diseases Grant 02971 to L.P.J. and National Institute of General Medical Sciences Grant 58884 to J.A.H.

¹ Abbreviations used are: NAPQI, N-acetyl-p-benzoquinone imine; GSH, glutathione; IL, interleukin; TNF- α , tumor necrosis factor- α ; NO, nitric oxide; iNOS, inducible nitric oxide synthase; IFN-y, interferon-y; MIF, macrophage migration inhibitory factor; MPT, mitochondrial permeability transition.

Several recent excellent reviews on acetaminophen toxicity have been recently published (Bessems and Vermeulen, 2001; Nelson et al., 2002). The following manuscript will review the role of covalent binding in toxicity as well as other factors recently identified that contribute to the toxicity. These factors include oxidative stress, nitrotyrosine formation, inflammatory cytokines, and the possible importance of mitochondrial permeability transition.

Metabolic Activation of Acetaminophen

Work from Dr. Gillette's laboratory firmly established the importance of metabolism in acetaminophen toxicity. It was shown that acetaminophen is metabolically activated by cytochrome P450 to form a reactive metabolite that covalently binds to protein (Mitchell et al., 1973). The reactive metabolite was found to be *N*-acetyl-*p*-benzoquinone imine (NAPQI¹), which is formed by a direct two-electron oxidation (Dahlin et al., 1984). More recently, the cytochromes 2E1, 1A2, 3A4, and 2A6 (Patten et al., 1993; Thummel et al., 1993; Chen et al., 1998) have been reported to oxidize acetaminophen to the reactive metabolite. Also, Dr. Gillette's laboratory showed that NAPQI is detoxified by glutathione (GSH) to form an acetaminophen-GSH conjugate. After a toxic dose of acetaminophen, total hepatic GSH is depleted by as much as 90%, and as a result, the metabolite

Address correspondence to: Dr. Laura James, Department of Pediatrics, Arkansas Children's Hospital, 800 Marshall Street, Little Rock, AR 72202. E-mail: jameslaurap@uams.edu

DRUG METABOLISM AND DISPOSITION

Aspet



FIG. 1. Schematic representation depicting the role of metabolism in acetaminophen toxicity.

covalently binds to cysteine groups on protein, forming acetaminophen-protein adducts (Mitchell et al., 1973). This mechanism is shown in Fig. 1.

To better understand the role of specific adducts in acetaminophen toxicity, our laboratory and Cohen's laboratory (Bartolone et al., 1988, 1989; Cohen and Khairallah, 1997) developed immunochemical assays that were very specific for acetaminophen covalently bound to cysteine groups on protein (Roberts et al., 1991). Using this antiserum, we reported the relationship between acetaminophen covalent binding and toxicity in mice in a competitive ELISA to quantify the adducts (Pumford et al., 1989, 1990a,b). Western blot assays were used to determine the molecular nature of the acetaminophen-protein adducts (Bartolone et al., 1988; Pumford et al., 1990a,b) and immunohistochemical analyses were used to determine the specific cells that contained the adducts (Bartolone et al., 1989; Roberts et al., 1991). These studies have been reviewed (Hinson et al., 1995; Pumford et al., 1997; Cohen and Khairallah, 1997).

Western blot assays for acetaminophen protein adducts have also been used to study toxicity in human overdose patients. In these studies, adducts have been detected in the blood of human overdose patients with severe toxicity (hepatic transaminases above 5000 IU/l) (James et al., 2001). Using high performance liquid chromatography with electrochemical detection, we recently reported the presence of adducts in the blood samples of overdose patients with mild to moderate toxicity, suggesting that the newer method may be a more sensitive analytical tool than Western blot for the detection of adducts in human sera (Muldrew et al., 2002). This assay may assist in the diagnosis of patients with acute liver failure of indeterminate etiology for which acetaminophen is suspect.

Biochemical Mechanisms of Toxicity

Events that produce hepatocellular death following the formation of acetaminophen protein adducts are poorly understood. One possible mechanism of cell death is that covalent binding to critical cellular proteins results in subsequent loss of activity or function and eventual cell death and lysis. Primary cellular targets have been postulated to be mitochondrial proteins, with resulting loss of energy production, as well as proteins involved in cellular ion control (Nelson, 1990). Tirmenstein and Nelson (1989) and Tsokos-Kuhn et al. (1988) reported alterations of plasma membrane ATPase activity following toxic doses of acetaminophen.

A number of proteins bound to acetaminophen have been isolated and identified. A review of the individual proteins that were isolated and identified by individual analysis has been published (Cohen and Khairallah, 1997; Pumford et al., 1997). Subsequently, Qui et al. (1998) used matrix-assisted laser desorption ionization mass spectrometry to identify 20 additional proteins containing covalently bound acetaminophen. Table 1 shows the proteins that have been identified to form adducts with acetaminophen. The proteins identified by our laboratory (Pumford et al., 1997) included a 100-kDa cytosolic protein determined by sequence analysis to be the enzyme, N-10-formyltetrahydrofolate dehydrogenase. This enzyme is involved in 1-carbon metabolism and oxidizes formaldehyde to carbon dioxide. Studies in mice reported that the activity of this enzyme was decreased by 20% at 2 h after toxic doses of acetaminophen (Pumford et al., 1997). A second protein, a 50-kDa mitochondrial protein, was isolated and determined to be glutamate dehydrogenase (Halmes et al., 1996). This enzyme reversibly metabolizes glutamate to α -ketoglutarate and ammonia. The activity of this enzyme was also decreased approximately 25% at 2 h. Based on the enzyme inhibition data for these two enzymes, it appeared that covalent binding with acetaminophen resulted in only partial inhibition of enzyme activities under toxic conditions in mice. Although it is plausible that partial inhibition of a large number of enzymes may contribute to cell death, the data generated questions relative to the validity of the hypothesis that covalent binding to critical proteins is the only mechanism of acetaminophen toxicity.

Loss of mitochondrial or nuclear ion balance has also been suggested to be a toxic mechanism involved in acetaminophen-mediated cell death since either of these losses can lead to increases in cytosolic

ACETAMINOPHEN-INDUCED HEPATOTOXICITY

TABLE 1

. . . 1.

Acetaminophen-protein adducts		
Mass	Fraction	Protein
kDa		
Proteins identified by isolation		
and sequence analysis ^a		
44	Microsomes	Glutamine synthetase
50	Mitochondria	Glutamate dehydrogenase
54	Mitochondria	Aldehyde dehydrogenase
55–58	Cytosol	Selenium (acetaminophen) binding protein
74–75	Nucleus	Lamin-A
100	Cytosol	N-10 Formyl THF dehydrogenase
Proteins identified by mass		
spectrometry ^b		
22	Cytosol, mitochondria	Glutathione peroxidase
28	Mitochondria	House keeping protein
29	Cytosol, microsomes	Thioether S-methyltransferase
16	Cytosol	Aryl sulfotransferase
32	Not known	Pyrophosphatase
29	Cytoskeleton	Tropomyosin 5
22	Cytosol, mitochondria	GSH peroxidase
52	Cytosol	Selenium binding protein
52	Cytosol	Acetaminophen binding protein
28	Cytosol	Proteasome subunit C8
45	Cytosol	Methionine adenosyl transferase
46	Ribosomes	Protein synthesis initiation factor 4A
56	Mitochondria	Aldehyde dehydrogenase
54	Cytosol	Aldehyde dehydrogenase
59	Mitochondria	ATP synthetase α -subunit
29	Cytosol	Carbonic anhydrase III
35	Peroxisomes	Urate oxidase
36	Mitochondria, peroxisome	2,4-Dienoyl-CoA reductase
22	Cytosol (macrophages)	Osteoblast-specific factor 3
23	Cytosol, mitochondria	Glutathione transferase π
40	Cytosol	Sorbitol dehydrogenase precursor
32	Cytosol	Glycine N-methyltransferase
32	Cytosol	3-Hydroxyanthraniliate 3,4-dioxygenase

THF, tetrahydrofuran.

Cohen et al. (1997).

^b Qiu et al. (1998).

Ca²⁺ concentrations, mitochondrial Ca²⁺ cycling, activation of proteases and endonucleases, and DNA strand breaks (Nelson, 1990; Ray et al., 1993; Salas and Corcoran, 1997). The effect of the addition of NAPQI on isolated mitochondria has been reported (Weis et al., 1992) and inhibition of mitochondrial respiration has been investigated as an important mechanism in acetaminophen toxicity (Donnelly et al., 1994).

Early Research on Oxidative Stress

Oxidative stress is another mechanism that has been postulated to be important in the development of acetaminophen toxicity. Thus, increased formation of superoxide would lead to hydrogen peroxide and peroxidation reactions by Fenton-type mechanisms. It has been shown that NAPQI reacts very rapidly with GSH ($k_1 = 3.2 \times 10^4$ M^{-1} s⁻¹ at pH 7.0) (Coles et al., 1988), and there are a number of potential mechanisms that have been suggested to play a role. Under conditions of NAPQI formation following toxic acetaminophen doses, GSH concentrations may be very low in the centrilobular cells, and the major peroxide detoxification enzyme, GSH peroxidase, which functions very inefficiently under conditions of GSH depletion (Nakamura et al., 1974), is expected to be inhibited. In addition, during formation of NAPQI by cytochrome P450, the superoxide anion is formed, with dismutation leading to hydrogen peroxide formation (Dai and Cederbaum, 1995). Also, others have suggested that peroxidation of acetaminophen to the semiquinone free radical would lead to redox cycling between the acetaminophen and the semiquinone. This mechanism may lead to increased superoxide and toxicity (de Vries,

1981). However, we found that the semiguinone reacted rapidly to form polymers and no evidence for reaction of oxygen was observed (Potter et al., 1989).

A significant amount of evidence has pointed to the potential involvement of oxidative stress in acetaminophen toxicity. Nakae et al. (1990) reported that administration of encapsulated superoxide dismutase decreased the toxicity of acetaminophen in the rat. Moreover, the iron chelator, deferoxamine, has been shown to decrease toxicity in rats (Sakaida et al., 1995). Our laboratory showed that deferoxamine caused a delay in the rate of development of acetaminophen toxicity in mice, but after 24 h, the relative amount of toxicity was not affected (Schnellmann et al., 1999). These data suggest that an iron-catalyzed Haber-Weiss reaction may also play a role in the development of oxidant stress and injury.

Role of Kupffer Cells

Several laboratories have studied the role of macrophage activation (Laskin et al., 1995) in acetaminophen toxicity. Kupffer cells are the phagocytic macrophages of the liver. When activated, Kupffer cells release numerous signaling molecules, including hydrolytic enzymes, eicosanoids, nitric oxide, and superoxide. Kupffer cells may also release a number of inflammatory cytokines, including IL-1, IL-6, and TNF- α (Laskin et al., 1995), and multiple cytokines are released in acetaminophen toxicity (Blazka et al., 1995; Hogaboam et al., 1999a, 2000; Bourdi et al., 2002a,b). Laskin et al. (1995) examined the role of Kupffer cells in acetaminophen hepatotoxicity by pretreating rats with compounds that suppress Kupffer cell function (gadolinium Downloaded from dmd.aspetjournals.org by guest on February 16, 2012



aspet



FIG. 2. Immunohistochemical detection of nitrotyrosine in an acetaminophen-treated mouse.

A, mouse treated with 300 mg/kg acetaminophen and sacrificed at 4 h after acetaminophen ($40\times$). The brown stain represents nitrotyrosine. B, mouse treated with saline and sacrificed at 4 h and stained for nitrotyrosine. ($40\times$). C, mouse treated with 300 mg/kg acetaminophen and sacrificed at 4 h after acetaminophen ($20\times$). D, mouse treated with saline and sacrificed at 4 h and stained for nitrotyrosine ($20\times$). D, mouse treated with saline and sacrificed at 4 h and stained for nitrotyrosine ($20\times$).

chloride and dextran sulfate). These investigators reported that rats pretreated with these compounds were less sensitive to the toxic effects of acetaminophen. Similar effects were reported in the mouse (Blazka et al., 1995). Goldin et al. (1996) showed that treatment of mice with liposomes containing dichloromethylene disphosphonate to knock out the Kupffer cells also decreased acetaminophen toxicity. Similarly, we reported that pretreatment of mice with gadolinium chloride or dextran sulfate decreased the toxic effects of acetaminophen (Michael et al., 1999). These studies suggested a critical role for Kupffer cells in the development of acetaminophen hepatotoxicity; however, recent work by Ju et al. (2002) came to a different conclusion. They pretreated mice with gadolinium chloride and found that the number of Kupffer cells in the liver was only partially decreased. Consistent with previous reports, they showed decreased acetaminophen toxicity in the pretreated mice. However, treatment of mice with dichloromethylene diphosphonate completely depleted the liver of Kupffer cells, but toxicity was increased. These data raise questions relative to the importance of Kupffer cells in acetaminophen toxicity.

Nitrotyrosine and Acetaminophen Toxicity

We recently showed that nitrotyrosine occurs in the centrilobular cells of the liver of acetaminophen-treated mice (Hinson et al., 1998). Immunohistochemical analysis of liver of acetaminophen-treated mice indicated that nitration occurred in the same cells that contained acetaminophen adducts and developed necrosis (Fig. 2) (Hinson et al.,

1998). We postulated that nitrotyrosine was formed by a reaction of peroxynitrite with tyrosine. Nitration of tyrosine has been shown to be an excellent biomarker of peroxynitrite formation (Kaur and Halliwell, 1994). It is formed by a rapid reaction between nitric oxide and superoxide, and we found that NO synthesis (serum levels of nitrate plus nitrite) was increased in acetaminophen toxicity (Hinson et al., 1998). Also, others reported the induction of hepatic iNOS in the acetaminophen-treated rat (Gardner et al., 1998). NO and superoxide react to produce peroxynitrite (ONOO⁻) at a rate near the diffusioncontrolled limit $(7 \times 10^9 \text{ M}^{-1} \text{ s}^{-1})$. Peroxynitrite is a species that not only leads to the nitration of tyrosine but is also a potent oxidant that can attack a wide range of biological targets (Pryor and Squadrito, 1995), and under conditions of reduced cellular oxidant scavenging capability, peroxynitrite is highly toxic (Beckman and Koppenol, 1996). The anion is relatively stable, but the acid rearranges to nitrate with a half-life of 1.9 s at pH 7.4; the pK is 7.5 (Beckman and Koppenol, 1996). Oxidation reactions can involve one or two-electron processes. Oxidation of lipids (Rubbo, 1998), proteins, or DNA bases (Pryor and Squadrito, 1995) has been reported. Thus, peroxynitrite is an excellent candidate for a toxic species. Moreover, it is normally detoxified by GSH/GSH peroxidase, and GSH is depleted in acetaminophen toxicity (Radi et al., 1991). GSH peroxidase is a key enzyme in this defense mechanism (Sies et al., 1997). Thus, a normal detoxification mechanism for peroxynitrite is impaired. Also, even though acetaminophen itself will detoxify peroxynitrite (Whiteman et al., 1996), the drug is metabolized rapidly in the mouse and concentrations are low at the time when nitration is observed. However, mechanisms other than peroxynitrite have been recently reported to lead to nitrotyrosine. Thus, myeloperoxidase has been shown to oxidize the nitrite ion to the NO₂ radical, and this is a nitrating species (Davis et al., 2001). Since myeloperoxidase is present in neutrophils and neutrophil recruitment is a late event in acetaminophen toxicity, this mechanism may not be important (Lawson et al., 2000). Also, nitrite may be oxidized by heme or free metals, leading to the NO₂ radical (Thomas et al., 2002).

Nitration of Tyrosine versus Oxidative Stress

The role of tyrosine nitration in acetaminophen toxicity was investigated using inhibitors of nitric oxide synthetase and mice that were genetically deficient in iNOS (Michael et al., 2001). A comparison of the time course for development of acetaminophen toxicity in iNOS knockout mice and wild-type mice indicated no difference in the histological development of toxicity between the two groups of mice. Serum levels of nitrate plus nitrite (a measure of NO synthesis) were increased in the wild-type mice but not in the iNOS knockout mice, and analysis of liver homogenates for tyrosine nitration indicated significant levels in the wild-type mice and a decreased amount in the knockout mice, possibly as a result of endothelial NOS. It was concluded that the major murine enzyme important in tyrosine nitration under normal wild-type conditions was iNOS. Interestingly, these experiments showed that acetaminophen caused a 3-fold increase in lipid peroxidation (oxidative stress) in the iNOS knockout mice but no increase in the wild-type mice. Thus, in the wild-type mice, toxicity was accompanied by tyrosine nitration, whereas in the iNOS knockout mice, toxicity was accompanied by oxidative stress. In a subsequent study (Hinson et al., 2002), we determined the effect of NOS inhibitors on acetaminophen toxicity in mice. The iNOS inhibitor aminoguanidine did not significantly alter acetaminophen toxicity, but it greatly decreased tyrosine nitration and increased lipid peroxidation. Thus, the iNOS inhibitor gave information similar to that obtained in the iNOS knockout mice. These data were interpreted to indicate that excess levels of superoxide were formed in acetaminophen toxicity. When NO is present, as in the wild-type mice, the superoxide preferentially reacts to form peroxynitrite, which nitrates proteins. In the absence of NO, superoxide leads to lipid peroxidation (Fig. 3). These data indicate the importance of NO in the disposition of superoxide, leading to oxidative stress.

Cytokines and Other Inflammatory Mediators in Acetaminophen Toxicity

Several laboratories have reported that inflammatory cytokines are increased in acetaminophen toxicity. Blazka reported the up-regulation of TNF- α and IL-1 α in the acetaminophen-treated mouse (Blazka et al., 1995, 1996). In addition, he showed that selective immunoneutralization of either TNF- α or IL-1 α partially decreased toxicity for a period of time. In vitro studies by Kuo showed that IL-1 β caused a dose-dependent up-regulation of NO in rat hepatocytes exposed to acetaminophen (Kuo et al., 1997). Consistent with these data, we recently found that IL-1 β and NO up-regulation in the acetaminophen-treated mouse are temporally related (James et al., 2003b). Although others have confirmed the up-regulation of TNF- α in acetaminophen toxicity (Simpson et al., 2000), its role in the mediation of toxicity is somewhat controversial. TNF- α knockout mice were not protected from toxicity (Boess et al., 1998), and a recent report showed that immunoneutralization of TNF- α had no effect on toxicity (Simpson et al., 2000). Studies in IL-10 knockout mice suggest that

OXIDATIVE TOXICITY



FIG. 3. Schematic representation depicting the role of oxidative stress in acetaminophen toxicity.

IL-10 is protective in APAP toxicity by controlling NO and iNOS formation (Bourdi et al., 2002a). In addition, the increased susceptibility of IL-10 knockout mice to acetaminophen was associated with increased levels of TNF- α and IFN- γ . Thus, it may be that proinflammatory cytokines contribute to the toxicity, and that they are regulated by anti-inflammatory cytokines, such as IL-10 and others.

Macrophage migration inhibitory factor (MIF) is a recently described protein that has characteristics of a cytokine, a hormone, and an enzyme. It may promote the up-regulation of other pro-inflammatory cytokines, adhesion molecules, matrix metalloproteinase-2 expression, NO release, and cyclooxygenase-2 and has counter-regulatory effects on endogenous glucocorticoids (Baugh and Bucala, 2002). Pohl and colleagues reported that MIF was up-regulated in acetaminophen toxicity (Bourdi et al., 2002b). Moreover, MIF knockout mice were less susceptible to acetaminophen and had decreased IFN- γ production and increased heat shock protein expression (Bourdi et al., 2002b). Heat shock proteins are considered to be a protective mechanism against various physiologic and environmental stressors (Bourdi et al., 2002b). In addition, cyclooxygenase-2 has been reported to be hepatoprotective in acetaminophen, possibly due to up-regulation of heat shock proteins (Reilly et al., 2001).

Various chemokines have also been reported to be up-regulated in acetaminophen toxicity. Chemokines were initially recognized for their role in leukocyte recruitment. However, Lawson et al. (2000) showed that neutrophils followed the development of toxicity and that administration of antibodies to B2 integrins had no effect on toxicity. Chemokine up-regulation in acetaminophen toxicity appears to be hepatoprotective. Hogaboam showed that macrophage inhibitory protein-2 (MIP-2) and other related chemokines promoted hepatocyte regeneration in acetaminophen-treated mice (Hogaboam et al., 1999a). In additional studies, viral vector delivery of MIP-2 to acetaminophen-treated mice was protective against acetaminophen toxicity (Hogaboam et al., 1999b). Studies in mice deficient in CCR2, the primary receptor for the chemokine MCP-1 (monocyte chemoattractant protein-1), showed these mice to have increased toxicity to acetaminophen and increased formation of TNF- α and IFN- γ (Hogaboam et al., 2000). Cumulatively, these studies suggest that cytokines play complex roles in the toxicity and that alterations in the equilibrium of pro- and anti-inflammatory cytokine formation contribute to the toxicity. Chemokines appear to facilitate hepatocyte regeneration following toxicity, likely by increasing the nuclear translocation of growth-regulatory transcription factors.

Superoxide Formation and Mitochondrial Dysfunction in Acetaminophen Toxicity

Superoxide may be formed via a number of mechanisms including formation from cytochrome P450 (Puntarulo and Cederbaum, 1996) and other enzymes. We evaluated the importance of activation of Kupffer cells, macrophages, or neutrophils (the so-called respiratory burst) in acetaminophen toxicity. This sudden excess utilization of oxygen by activated phagocytes is a result of increased activity of the enzyme, NADPH-oxidase. The result is release of superoxide anion at the outer surface of the plasma membrane (Baggiolini and Wymann, 1990). We utilized mice genetically deficient in Gp91 phox, a critical subunit of NADPH oxidase (James et al., 2003a), and found that these mice and wild-type mice had comparable toxicity to acetaminophen. Also, both knockout mice and wild-type mice had the same degree of tyrosine nitration in the liver. An analysis of mitochondrial glutathione disulfide was found to be comparable in the two groups (James et al., 2003a). These data suggest that the increased superoxide in acetaminophen toxicity is not from activated macrophages.

Available data suggest that mitochondrial dysfunction may be an important mechanism in acetaminophen-induced hepatotoxicity. It is known that mitochondrial permeability transition (MPT) occurs with formation of superoxide, and this may be the source of superoxide leading to peroxynitrite and tyrosine nitration. MPT represents an abrupt increase in the permeability of the inner mitochondrial membrane to small molecular weight solutes. Oxidants such as peroxides and peroxynitrite, Ca²⁺, and P_i promote the onset of MPT, whereas Mg²⁺, ADP, low pH, and high membrane potential oppose onset. Associated with the permeability change is membrane depolarization, uncoupling of oxidative phosphorylation, release of intramitochondrial ions and metabolic intermediates, and mitochondrial swelling. MPT mechanisms have been reviewed by Lemasters et al. (1998). In the late 1980s it was shown that cyclosporin A specifically blocked the onset of MPT in a saturable manner, implying a protein channel or pore. It has been subsequently recognized that this pore transports both anionic and cationic solutes of mass less than 1500 Da and is the previously identified multiple conductance channel. Of particular note is the fact that oxidation of vicinal thiols in the pore promote an open conductance state (Lemasters et al., 1998). Moldeus and Orrenius' laboratory reported that addition of NAPQI to isolated rat liver mitochondria caused a decrease in synthesis of ATP and an increase in release of sequestered Ca²⁺. This release was blocked by cyclosporin A (Weis et al., 1992). These data are consistent with the hypothesis that NAPQI causes MPT, as has been reported about other quinones (Palmeira and Wallace, 1997). This is presumably a result of NAPOImediated oxidation of the vicinal thiols at the MPT pore. NAPQI is known to be both an oxidizing agent and an arylating agent, and Tirmenstein and Nelson (1989) have reported that acetaminophen leads to oxidation of protein thiols. Also, McLean and coworkers have reported that inhibitors of MPT decrease acetaminophen toxicity in rat liver slices and in vivo (Beales and McLean, 1996). Lastly, Grewal reported that late addition of N-acetylcysteine to freshly isolated mouse liver hepatocytes had no effect on the toxicity; however, dithiothreitol decreased the further development of toxicity (Grewal and Racz, 1993). Also, in similar experiments with isolated hamster hepatocytes, the late addition of dithiothreitol to cells washed free of acetaminophen prevented the further decrease in cell viability and appeared to reverse morphological changes (Tee et al., 1986). Dithiothreitol is known to reduce the vicinal thiols at the MPT pore and

MITOCHONRIAL PERMEABILITY TRANSITION (MPT) IN APAP HEPATOTOXICITY



FIG. 4. Schematic representation depicting the role of mitochondrial permeability transition in acetaminophen toxicity.

prevent the further progression of toxicity (Palmeira and Wallace, 1997). These data are consistent with a mechanism whereby acetaminophen toxicity is by NAPQI-mediated oxidation of thiol groups in mitochondria leading to MPT. The toxicity is mediated by mitochondrial dysfunction resulting in production of reactive oxygen/ nitrogen species (Fig. 4).

Summary

Dr. Gillette and researchers in his laboratory made fundamental contributions to our understanding of the pathogenesis of drug and chemical toxicity. With the analgesic acetaminophen it was shown that the hepatic necrosis was not a result of the parent compound. Metabolism by cytochrome P450 enzymes to a reactive metabolite was necessary for the production of toxicity. After therapeutic doses of acetaminophen, the reactive metabolite was efficiently detoxified by conjugation with GSH. However, after toxic doses, GSH was depleted by the conjugation reaction and the metabolite covalently bound to protein. Covalent binding correlated with development of toxicity. Treatment with cysteine to increase GSH detoxification was shown to be an effective antidote against toxicity. The latter finding led to the development of N-acetylcysteine as the current clinical antidote. Since these initial findings, many investigators have added additional details relative to the concept of how acetaminophen is toxic, and some of these are discussed in this review. Despite these recent studies, the concept of metabolic activation as the critical initiating event in the development of toxicity has remained. Moreover, metabolic activation has proven to be a general mechanism relevant for the development of toxicity for many drugs and chemicals. Dr. Gillette reviewed his major contributions to science in the Annual Review of Pharmacology and Toxicology (Gillette, 2000) and in Drug Metabolism Reviews (Gillette, 1995).

References

- Baggiolini M and Wymann MP (1990) Turning on the respiratory burst. *Trends Biochem Sci* 15:69–72.
- Bartolone JB, Birge RB, Sparks K, Cohen SD, and Khairallah EA (1988) Immunochemical analysis of acetaminophen covalent binding to proteins. Partial characterization of the major acetaminophen-binding liver proteins. *Biochem Pharmacol* 37:4763–4774.
- Bartolone JB, Cohen SD, and Khairallah EA (1989) Immunohistochemical localization of acetaminophen-bound liver proteins. *Fundam Appl Toxicol* 13:859–862.
- Baugh JA and Bucala R (2002) Macrophage migration inhibitory factor. Crit Care Med 30 (Suppl):S27–S35.
- Beales D and McLean AE (1996) Protection in the late stages of paracetamol-induced liver cell injury with fractose, cyclosporin A and trifluoperazine. *Toxicology* **107**:201–208.
- Beckman JS and Koppenol WH (1996) Nitric oxide, superoxide, and peroxynitrite: the good, the bad and ugly. Am J Physiol 271 (Pt 1):C1424-C1437.
 Becomes Level Variation and Colling and Coll
- Bessems JG and Vermeulen NP (2001) Paracetamol (acetaminophen)-induced toxicity: molecular and biochemical mechanisms, analogues and protective approaches. *Crit Rev Toxicol* 31:55–138.

METABOLISM AND DISPOSIT

spet

 \square

JG METABOLISM AND DISPOSITIC

spet

- Blazka ME, Elwell MR, Holladay SD, Wilson RE, and Luster MI (1996) Histopathology of acetaminophen-induced liver changes: role of interleukin 1 alpha and tumor necrosis factor resp
- alpha. Toxicol Pathol 24:181–189.
 Blazka ME, Wilmer JL, Holladay SD, Wilson RE, and Luster MI (1995) Role of proinflammatory cytokines in acetaminophen hepatotoxicity. Toxicol Appl Pharmacol 133:43–52.
- Boess F, Bopst M, Althaus R, Polsky S, Cohen SD, Eugster HP, and Boelsterli UA (1998) Acetaminophen hepatotoxicity in tumor necrosis factor/lymphotoxin-alpha gene knockout mice. *Hepatology* 27:1021–1029.
- Bourdi M, Masubuchi Y, Reilly TP, Amouzadeh HR, Martin JL, George JW, Shah AG, and Pohl LR (2002a) Protection against acetaminophen-induced liver injury and lethality by interleukin 10: role of inducible nitric oxide synthase. *Hepatology* 35:289–298.
- Bourdi M, Reilly TP, Elkahloun AG, George JW, and Pohl LR (2002b) Macrophage migration inhibitory factor in drug-induced liver injury: a role in susceptibility and stress responsiveness. *Biochem Biophys Res Commun* 294:225–230.
- Chen W, Koenigs LL, Thompson SJ, Peter RM, Rettie AE, Trager WF, and Nelson SD (1998) Oxidation of acetaminophen to its toxic quinone imine and nontoxic catechol metabolites by baculovirus-expressed and purified human cytochromes P450 2E1 and 2A6. *Chem Res Toxicol* 11:295–301.
- Cohen SD and Khairallah EA (1997) Selective protein arylation and acetaminophen-induced hepatotoxicity. *Drug Metab Rev* **29:**59–77.
- Cohen SD, Pumford NR, Khairallah EA, Boekelheide K, Pohl LR, Amouazdeh HR, and Hinson JA (1997) Contemporary issues in toxicology: selective protein covalent binding and target organ toxicity. *Toxicol Appl Pharmacol* 143:1–12.
- Coles B, Wilson I, Wardman P, Hinson JA, Nelson SD, and Ketterer B (1988) The spontaneous and enzymatic reaction of N-acetyl-p-benzoquinonimine with glutathione: a stopped-flow kinetic study. Arch Biochem Biophys 264:253–260.
- Dahlin DC, Miwa GT, Lu AY, and Nelson SD (1984) N-Acetyl-p-benzoquinone imine: a cytochrome P-450-mediated oxidation product of acetaminophen. *Proc Natl Acad Sci USA* 81:1327–1331.
- Dai Y and Cederbaum AI (1995) Cytotoxicity of acetaminophen in human cytochrome P4502E1transfected HepG2 cells. J Pharmacol Exp Ther 273:1497–1505.
- Davis KL, Martin E, Turko IV, and Murad F (2001) Novel effects of nitric oxide. Annu Rev Pharmacol Toxicol 41:203–236.
- de Vries J (1981) Hepatotoxic metabolic activation of paracetamol and its derivatives phenacetin and benorilate: oxygenation or electron transfer? *Biochem Pharmacol* 30:399–402.
- Donnelly PJ, Walker RM, and Racz WJ (1994) Inhibition of mitochondrial respiration in vivo is an early event in acetaminophen-induced hepatotoxicity. Arch Toxicol 68:110–118.
- Gardner CR, Heck DE, Yang CS, Thomas PE, Zhang XJ, DeGeorge GL, Laskin JD, and Laskin DL (1998) Role of nitric oxide in acetaminophen-induced hepatotoxicity in the rat. *Hepatology* 27:748–754.
- Gillette JR (1995) Keynote address: man, mice, microsomes, metabolites and mathematics 40 years after the revolution. *Drug Metab Rev* 27:1–44.
- Gillette JR (2000) Laboratory of Chemical Pharmacology, National Heart, Lung and Blood Institute, NIH: a short history. Annu Rev Pharmacol Toxicol 40:18–41.
- Goldin RD, Ratnayaka ID, Breach CS, Brown IN, and Wickramasinghe SN (1996) Role of macrophages in acetaminophen (paracetamol)-induced hepatotoxicity. J Pathol 179:432–435. Grewal KK and Racz WJ (1993) Intracellular calcium disruption as a secondary event in acetaminophen-induced hepatotoxicity. Can J Physiol Pharmacol 71:26–33.
- Gujral JS, Knight TR, Farhood A, Bajt ML, and Jaeschke H (2002) Mode of cell death after acetaminophen overdose in mice: apoptosis or oncotic necrosis? *Toxicol Sci* 67:322–328. Halmes NC, Hinson JA, Martin BM, and Pumford NR (1996) Glutamate dehydrogenase
- rannes NC, Finson JA, Martin BM, and Funnou NK (1990) Outaniae denyarogenase covalently binds to a reactive metabolite of acetaminophen. *Chem Res Toxicol* 9:541–546. Hinson JA, Bucci TJ, Irwin LK, Michael SL, and Mayeux PR (2002) Effect of inhibitors of nitric
- oxide synthase on acetaminophen-induced hepatotoxicity in mice. *Nitric Oxide* 6:160-167. Hinson JA, Pike SL, Pumford NR, and Mayeux PR (1998) Nitrotyrosine-protein adducts in
- hepatic centrilobular areas following toxic doses of acetaminophen in mice. *Chem Res Toxicol* **11**:604–607. Hinson JA, Pumford NR, and Roberts DW (1995) Mechanisms of acetaminophen toxicity:
- immunochemical detection of drug-protein adducts. Drug Metab Rev 27:73–92.
- Hogaboam CM, Bone-Larson CL, Steinhauser ML, Lukacs NW, Colletti LM, Simpson KJ, Strieter RM, and Kunkel SL (1999a) Novel CXCR2-dependent liver regenerative qualities of ELR-containing CXC chemokines. *FASEB J* 13:1565–1574.
- Hogaboam CM, Bone-Larson CL, Steinhauser ML, Matsukawa A, Gosling J, Boring L, Charo IF, Simpson KJ, Lukacs NW, and Kunkel SL (2000) Exaggerated hepatic injury due to acetaminophen challenge in mice lacking C-C chemokine receptor 2. Am J Pathol 156:1245– 1252.
- Hogaboam CM, Simpson KJ, Chensue SW, Steinhauser ML, Lukacs NW, Gauldie J, Strieter RM, and Kunkel SL (1999b) Macrophage inflammatory protein-2 gene therapy attenuates adenovirus- and acetaminophen-mediated hepatic injury. *Gene Ther* 6:573–584. James LP, Farrar HC, Sullivan JE, Givens TG, Kearns GL, Wasserman GS, Walson PD, Hinson
- James LP, Farrar HC, Sullivan JE, Givens TG, Kearns GL, Wasserman GS, Walson PD, Hinson JA, Pumford NR and Pediatric Pharmacology Research Unit Network, NICHD (2001) Measurement of acetaminophen-protein adducts in children and adolescents with acetaminophen overdoses. J Clin Pharmacol 41:846–851.
- James LP, McCullough SS, Knight TR, Jaeschke H, and Hinson JA (2003a) Acetaminophen toxicity in mice lacking NADPH oxidase activity: role of peroxynitrite formation and mitochondrial oxidant stress. *Free Radic Res*, in press.
- James LP, McCullough SS, Lamps LW, and Hinson JA (2003b). Effect of N-acetylcysteine on acetaminophen toxicity in mice: relationship to reactive nitrogen and cytokine formation. *Toxicol Sci* 75:458–467.
- Ju C, Reilly TP, Bourdi M, Radonovich MF, Brady JN, George JW, and Pohl LR (2002) Protective role of Kupffer cells in acetaminophen-induced hepatic injury in mice. *Chem Res Toxicol* 15:1504–1513.
- Kaur H and Halliwell B (1994) Evidence for nitric oxide-mediated oxidative damage in chronic inflammation. Nitrotyrosine in serum and synovial fluid from rheumatoid patients. *FEBS Lett* 350:9–12.
- Kuo PC, Schroeder RA, and Loscalzo J (1997) Nitric oxide and acetaminophen-mediated oxidative injury: modulation of interleukin-1-induced nitric oxide synthesis in cultured rat hepatocytes. J Pharmacol Exp Ther 282:1072–1083.
- Laskin DL, Gardner CR, Price VF, and Jollow DJ (1995) Modulation of macrophage functioning abrogates the acute hepatotoxicity of acetaminophen. *Hepatology* 21:1045–1050.

- Lawson JA, Farhood A, Hopper RD, Bajt ML, and Jaeschke H (2000) The hepatic inflammatory response after acetaminophen overdose: role of neutrophils. *Toxicol Sci* 54:509–516.
- Lawson JA, Fisher MA, Simmons CA, Farhood A, and Jaeschke H (1999) Inhibition of Fas receptor (CD95)-induced caspase activation and apoptosis by acetaminophen in mice. *Toxicol Appl Pharmacol* 156:179–186.
- Lemasters JJ, Nieminen AL, Qian T, Trost LC, Elmore SP, Nishimura Y, Crowe RA, Cascio WE, Bradham CA, Brenner DA, and Herman B (1998) The mitochondrial permeability transition in cell death: a common mechanism in necrosis, apoptosis and autophagy. *Biochim Biophys Acta* 1366:177–196.
- Litovitz TL, Klein-Schwartz W, Rodgers GC Jr, Cobaugh DJ, Youniss J, Omslaer JC, May ME, Woolf AD, and Benson BE (2002) 2001 annual report of the American Association of Poison Control Centers Toxic Exposure Surveillance System. Am J Emerg Med 20:391–452.
- Michael SL, Mayeux PR, Bucci TJ, Warbritton AR, Irwin LK, Pumford NR, and Hinson JA (2001) Acetaminophen-induced hepatotoxicity in mice lacking inducible nitric oxide synthase activity. *Nitric Oxide* 5:432–441.
- Michael SL, Pumford NR, Mayeux PR, Niesman MR, and Hinson JA (1999) Pretreatment of mice with macrophage inactivators decreases acetaminophen hepatotoxicity and the formation of reactive oxygen and nitrogen species. *Hepatology* 30:186–195.
- Mitchell JR, Jollow DJ, Potter WZ, Gillette JR, and Brodie BB (1973) Acetaminophen-induced hepatic necrosis. IV. Protective role of glutathione. J Pharmacol Exp Ther 187:211–217.
- Muldrew KL, James LP, Coop L, McCullough SS, Hendrickson HP, Hinson JA, and Mayeux PR (2002) Determination of acetaminophen-protein adducts in mouse liver and serum and human serum after hepatotoxic doses of acetaminophen using high- performance liquid chromatography with electrochemical detection. *Drug Metab Dispos* 30:446–451.
- Nakae D, Yoshiji H, Yamamoto K, Maruyama H, Kinugasa T, Takashima Y, Denda A, and Konishi Y (1990) Influence of timing of administration of liposome-encapsulated superoxide dismutase on its prevention of acetaminophen-induced liver cell necrosis in rats. Acta Pathol Jpn 40:568–573.
- Nakamura W, Hosoda S, and Hayachi K (1974) Purification and properties of rat liver glutathione peroxidase. *Biochim Biophys Acta* 358:251–261.
- Nelson SD (1990) Molecular mechanisms of the hepatotoxicity caused by acetaminophen. Semin Liver Dis 10:267–278.
- Nelson SG, Wan Z, and Stan MA (2002) S(N)2 ring opening of beta-lactones: an alternative to catalytic asymmetric conjugate additions. J Org Chem 67:4680-4683.
- Palmeira CM and Wallace KB (1997) Benzoquinone inhibits the voltage-dependent induction of the mitochondrial permeability transition caused by redox-cycling naphthoquinones. *Toxicol Appl Pharmacol* 143:338–347.
- Patten CJ, Thomas PE, Guy RL, Lee M, Gonzalez FJ, Guengerich FP, and Yang CS (1993) Cytochrome P450 enzymes involved in acetaminophen activation by rat and human liver microsomes and their kinetics. *Chem Res Toxicol* 6:511–518.
- Potter DW, Pumford NR, Hinson JA, Benson RW, and Roberts DW (1989) Epitope characterization of acetaminophen bound to protein and nonprotein sulfhydryl groups by an enzymelinked immunosorbent assay. J Pharmacol Exp Ther 248:182–189.
- Prescott LF (1980) Hepatotoxicity of mild analgesics Br J Clin Pharmacol 10 (Suppl 2):373S– 379S.
- Pryor WA and Squadrito GL (1995) The chemistry of peroxynitrite: a product from the reaction of nitric oxide with superoxide. Am J Physiol 268(Pt 1):L699–L722.
- Pumford NR, Halmes NC, Martin BM, Cook RJ, Wagner C, and Hinson JA (1997) Covalent binding of acetaminophen to N-10-formyltetrahydrofolate dehydrogenase in mice. J Pharmacol Exp Ther 280:501–505.
- Pumford NR, Hinson JA, Benson RW, and Roberts DW (1990a) Immunoblot analysis of protein containing 3-(cystein-S-yl)acetaminophen adducts in serum and subcellular liver fractions from acetaminophen-treated mice. *Toxicol Appl Pharmacol* 104:521–532.
- Pumford NR, Hinson JA, Potter DW, Rowland KL, Benson RW, and Roberts DW (1989) Immunochemical quantitation of 3-(cystein-S-yl)acetaminophen adducts in serum and liver proteins of acetaminophen-treated mice. J Pharmacol Exp Ther 248:190–196.
- Pumford NR, Roberts DW, Benson RW, and Hinson JA (1990b) Immunochemical quantitation of 3-(cystein-S-yl)acetaminophen protein adducts in subcellular liver fractions following a hepatotoxic dose of acetaminophen. *Biochem Pharmacol* 40:573–579.
- Puntarulo S and Cederbaum AI (1996) Role of cytochrome P-450 in the stimulation of microsomal production of reactive oxygen species by ferritin. *Biochim Biophys Acta* 1289:238-246.
- Qiu Y, Benet LZ, and Burlingame AL (1998) Identification of the hepatic protein targets of reactive metabolites of acetaminophen in vivo in mice using two-dimensional gel electrophoresis and mass spectrometry. J Biol Chem 273:17940–17953.
- Radi R, Beckman JS, Bush KM, and Freeman BA (1991) Peroxynitrite oxidation of sulfhydryls. The cytotoxic potential of superoxide and nitric oxide. J Biol Chem 266:4244–4250.
- Ray SD, Kamendulis LM, Gurule MW, Yorkin RD, and Corcoran GB (1993) Ca2+ antagonists inhibit DNA fragmentation and toxic cell death induced by acetaminophen. FASEB J 7:453– 463.
- Reilly TP, Brady JN, Marchick MR, Bourdi M, George JW, Radonovich MF, Pise-Masison CA, and Pohl LR (2001) A protective role for cyclooxygenase-2 in drug-induced liver injury in mice. *Chem Res Toxicol* 14:1620–1628.
- Roberts DW, Bucci TJ, Benson RW, Warbritton AR, McRae TA, Pumford NR, and Hinson JA (1991) Immunohistochemical localization and quantification of the 3-(cystein-S-yl)acetaminophen protein adduct in acetaminophen hepatotoxicity. *Am J Pathol* 138:359– 371.
- Rubbo H (1998) Nitric oxide and peroxynitrite in lipid peroxidation. Medicina 58:361-366.
- Sakaida I, Kayano K, Wasaki S, Nagatomi A, Matsumura Y, and Okita K (1995) Protection against acetaminophen-induced liver injury in vivo by an iron chelator, deferoxamine. *Scand J Gastroenterol* 30:61–67.
- Salas VM and Corcoran GB (1997) Calcium-dependent DNA damage and adenosine 3',5'-cyclic monophosphate-independent glycogen phosphorylase activation in an in vitro model of acetaminophen-induced liver injury. *Hepatology* 25:1432–1438.
- Schnellmann JG, Pumford NR, Kusewitt DF, Bucci TJ, and Hinson JA (1999) Deferoxamine delays the development of the hepatotoxicity of acetaminophen in mice. *Toxicol Lett* 106: 79–88.
- Sies H, Sharov VS, Klotz LO, and Briviba K (1997) Glutathione peroxidase protects against peroxynitrite-mediated oxidations. A new function for selenoproteins as peroxynitrite reductase. J Biol Chem 272:27812–27817.

Downloaded from dmd.aspetjournals.org by guest on February 16,

2012

DRUG METABOLISM AND DISPOSITI

Gspet

1506

- Simpson KJ, Lukacs NW, McGregor AH, Harrison DJ, Strieter RM, and Kunkel SL (2000) Inhibition of tumour necrosis factor alpha does not prevent experimental paracetamol-induced hepatic necrosis. J Pathol 190:489–494.
- Tee LB, Boobis AR, Huggett AC, and Davies DS (1986) Reversal of acetaminophen toxicity in isolated hamster hepatocytes by dithiothreitol. *Toxicol Appl Pharmacol* 83:294–314.
- Thomas DD, Espey MG, Vitek MP, Miranda KM, and Wink DA (2002) Protein nitration is mediated by heme and free metals through Fenton-type chemistry: an alternative to the NO/O₂⁻ reaction. *Proc Natl Acad Sci USA* 99:12691–12696.
- Thummel KE, Lee CA, Kunze KL, Nelson SD, and Slattery JT (1993) Oxidation of acetaminophen to N-acetyl-p-aminobenzoquinone imine by human CYP3A4. *Biochem Pharmacol* 45:1563–1569.
- Tirmenstein MA and Nelson SD (1989) Subcellular binding and effects on calcium homeostasis produced by acetaminophen and a nonhepatotoxic regioisomer, 3'-hydroxyacetanilide, in mouse liver. J Biol Chem 264:9814–9819.
- Tsokos-Kuhn JO, Hughes H, Smith CV, and Mitchell JR (1988) Alkylation of the liver plasma membrane and inhibition of the Ca2+ ATPase by acetaminophen. *Biochem Pharmacol* 37:2125–2131.
- Weis M, Kass GE, Orrenius S, and Moldeus P (1992) N-Acetyl-p-benzoquinone imine induces Ca2+ release from mitochondria by stimulating pyridine nucleotide hydrolysis. J Biol Chem 267:804–809.
- Whiteman M, Kaur H, and Halliwell B (1996) Protection against peroxynitrite dependent tyrosine nitration and alpha 1-antiproteinase inactivation by some anti-inflammatory drugs and by the antibiotic tetracycline. Ann Rheum Dis 55:383–387.



Philip Mayeux received his Ph.D. in pharmacology from Tulane University in 1987. After completing postdoctoral fellowships with Dr. Perry V. Halushka at the Medical University of South Carolina and Dr. Sudhir V. Shah at Tulane University, he joined the faculty of the Department of Pharmacology and Toxicology at the University of Arkansas for Medical Sciences. He is an Associate Professor of Pharmacology and Toxicology and directs the Ph.D. program in Pharmacology. His research interests are on the role of reactive nitrogen species in renal and hepatic failure. He is on the editorial board of the

Journal of Pharmacology and Experimental Therapeutics and is a member of ASPET, SOT, American Society of Nephrology, and Nitric Oxide Society.



Jack Hinson received a Ph.D. in biochemistry from Vanderbilt University in 1972. Subsequently, he was a postdoctoral fellow and a senior staff fellow in the Laboratory of Chemical Pharmacology at the National Institutes of Health where he worked with Dr. James R. Gillette. In 1980 he transferred to FDA's National Center for Toxicological Research where he was Chief of the Biochemical Mechanisms Branch. In 1990 he assumed his current position of Professor and Director of the Division of Toxicology, Department of Pharmacology and Toxicology at the University of Arkansas for Medical Sciences. He directs a Ph.D. program in Inter-

disciplinary Toxicology. His research interests are on the role of drug metabolism and reactive oxygen/nitrogen species in drug-induced liver toxicity. He is the editor of *Drug Metabolism Reviews* and is on the editorial boards of *Toxicology and Applied Pharmacology* and the *Journal of Toxicology and Environmental Health.* He serves as a member of a National Institutes of Health advisory panel and is a member of SOT, ASPET, Nitric Oxide Society, and ISSX.



Laura James received her M.D. from the University of South Carolina in 1989. She completed an internship and residence in Pediatrics at University of Arkansas and Arkansas Children's Hospital in 1992. She then completed a fellowship in Pediatric Emergency Medicine at the University of Alabama at Birmingham in 1994, followed by a fellowship in Pediatric Pharmacology/Toxicology at University of Arkansas for Medical Sciences in 1996. She is an Associate Professor in the Department of Pediatrics at the University of Arkansas for Medical Sciences. Her research interests include acet-

aminophen toxicity and drug metabolism in chil-

dren. She is a member of the American Academy of Pediatrics, the American Society for Clinical Pharmacology and Therapeutics, The American College of Clinical Toxicology, and Society for Pediatric Research.